

## Medication Information Sheet

**azaCITIDine** (ay-za-SYE-ti-deen)

*This document provides general information about your medication. It does not replace the advice of your health care professional. Always discuss your therapy with your health care professional and refer to the package insert for more details.*

**Other Name:** Vidaza®

**Appearance:** White, milky suspension for injection

**What is this medication for?**

- For treating certain types of blood disorders, such as myelodysplastic syndromes (MDS) or acute myeloid leukemia (AML)

**What should I do before I have this medication?**

- Tell your doctor and pharmacist if you have/had significant medical condition(s), especially if you have / had heart problems (including abnormal heart rhythm), kidney, liver or lung problems, or any allergies.
- People who have cancer or leukemia are at a higher risk of developing other cancers/leukemias (usually some years later) or blood clots. Some cancer medications may increase these risks, especially if used for a prolonged period of time. You should discuss any concerns with your doctor.

**How will this medication affect sex, pregnancy and breastfeeding?**

- azaCITIDine can harm the unborn baby and should not be used by pregnant women.
- Do not use this drug if you are pregnant. If there is ANY chance that you or your partner may become pregnant, you and your partner together must: ► **Use 2 effective forms of birth control at the same time** while taking this drug: Keep using birth control until **6 months** after the last dose (general recommendation). Discuss with your healthcare team.
- Tell your doctor right away if you or your partner becomes pregnant.
- Do not breastfeed while taking this drug.
- Effects on Fertility: Probable

**How is this medication given?**

- This drug is given by injection under the skin.

## What else do I need to know while on this medication?

- This medication can interact with other medications and can result in the treatment not working as well or cause severe side effects.
- Make sure your health care team knows about all your medications (prescription, over-the-counter, herbals and supplements). Check with your health care team before starting or stopping any of them.
- For mild aches and pain:
  - You may take acetaminophen (Tylenol®) tablets. Ask your health care team about the right dose for you.
  - Talk to your health care team first before starting ibuprofen (Advil®, Motrin®), naproxen (Aleve®) or aspirin (acetylsalicylic acid, ASA), since these may increase your chance of bleeding.
  - If you are already taking aspirin regularly, such as for heart conditions; Do not stop it- talk to your health care team first.
  - If you feel unwell, take your temperature before using any of these drugs. They may hide a fever. **Phone your health care team or go to the closest emergency room right away if you have a fever.** See the [Fever](#) pamphlet for more information.

## What are the side effects of this medication?

Common side effects usually occur in more than 1 out of every 10 patients. Other side effects are less common, but may be severe. You may not have all of the side effects below. You may have side effects that are not listed.

Side effects and what to do	When to contact doctor?
<b>More Common Side Effects</b>	
<p><b><i>Nausea and vomiting</i></b></p> <ul style="list-style-type: none"> <li>• Drink clear fluids and avoid large meals. Get fresh air and rest.</li> <li>• Limit spicy, fried foods or foods with a strong smell.</li> <li>• Take anti-nausea drug(s) exactly as directed by your doctor. It is easier to prevent nausea than to treat it.</li> <li>• Contact your doctor if nausea lasts more than 48 hours or vomiting for more than 24 hours.</li> <li>• Also see Nausea &amp; Vomiting pamphlet.*</li> </ul>	<p>Contact your health care team if no improvement or if severe</p>

\*The most updated version and more symptom control information can be found on: <http://www.cancercare.on.ca/druginfo>  
 Prepared with input from the Cancer Care Ontario-Medication Information Sheets Working Group.  
 April 2016

Side effects and what to do	When to contact doctor?
<b>More Common Side Effects</b>	
<p><b>Unusual bleeding or bruising</b></p> <p>You may have black stools, cough up blood, blood in your urine, purple or red dots on your skin or bleeding that will not stop.</p> <p><b>Fever, chills, infection</b></p> <p>You have a fever if your temperature taken in your mouth (oral temperature) is:</p> <ul style="list-style-type: none"> <li>• <b>38.3°C (100.9°F) or higher at any time OR</b></li> <li>• <b>38.0°C (or 100.4°F) or higher for at least one hour.</b></li> </ul> <p>While you are getting chemotherapy treatments:</p> <ul style="list-style-type: none"> <li>• Keep a digital thermometer at home and take your temperature if you feel hot or unwell (for example, chills).</li> <li>• Avoid taking medications that treat a fever before you take your temperature (for example, Tylenol®, acetaminophen, Advil® or ibuprofen) as they may hide a fever.</li> <li>• Do not eat or drink anything hot or cold right before taking your temperature.</li> <li>• Wash your hands often.</li> <li>• Check with your doctor before getting any vaccines, surgeries or visiting your dentist.</li> </ul> <p><b>If you have a fever, talk to your health care team or go to the closest emergency room.</b></p> <p>See our <a href="#">Neutropenia (Low white blood cell count)</a> pamphlet for more information.</p>	<p>Get emergency medical help right away</p>
<p><b>Constipation</b></p> <ul style="list-style-type: none"> <li>• Eat a balanced diet with fibres such whole grains, fruit and raw vegetables.</li> <li>• Drink plenty of fluids. Try light exercise regularly.</li> <li>• Speak to your doctor if no bowel movement for 3 or more days.</li> <li>• Also see Constipation Pamphlet.*</li> </ul>	<p>Contact your health care team if no improvement or if severe</p>
<p><b>Reactions at the injection site</b> (redness, itchiness, bruising, mild rash or swelling)</p>	<p>Contact your health care team if no improvement or if severe</p>

\*The most updated version and more symptom control information can be found on: <http://www.cancercare.on.ca/druginfo>  
 Prepared with input from the Cancer Care Ontario-Medication Information Sheets Working Group.  
 April 2016

Side effects and what to do	When to contact doctor?
<b>More Common Side Effects</b>	
<p><b>Diarrhea</b></p> <ul style="list-style-type: none"> <li>• Drink plenty of clear fluids. Limit hot, spicy, fried foods, foods/drinks with caffeine, orange or prune juice. Try a low-fiber BRAT diet (Bananas, white Rice, Apple sauce, Toast made with white bread).</li> <li>• Take anti-diarrhea drug(s) if given to you by your doctor.</li> <li>• Also see Diarrhea pamphlet.*</li> </ul>	Contact your health care team if no improvement or if severe
<p><b>Tiredness</b></p> <ul style="list-style-type: none"> <li>• Rest often; take naps if needed. Move slowly when getting up.</li> <li>• Eat well-balanced meals and drink plenty of fluids. Light exercise may help.</li> <li>• Do not drive a motor vehicle or operate machinery when feeling tired.</li> </ul>	Contact your health care team if no improvement or if severe
<p><b>Headache; mild joint or muscle pain</b></p> <ul style="list-style-type: none"> <li>• Take painkiller(s) as directed, if given to you by your doctor.</li> <li>• Otherwise, take acetaminophen (Tylenol®) tablets as needed for mild aches and pains. Ask your doctor or pharmacist about the right dose for you.</li> <li>• Rest often, but may try light exercise.</li> </ul>	Contact your health care team if no improvement or if severe
<p><b>Poor Appetite; don't feel like eating; weight loss</b></p> <ul style="list-style-type: none"> <li>• Eat foods that you like and try to eat regular small meals.</li> <li>• Use meal supplements if possible. See a dietitian.</li> </ul>	Contact your health care team if no improvement or if severe
<p><b>Dizziness, lightheadedness</b></p> <ul style="list-style-type: none"> <li>• Do not drive a motor vehicle or operate machinery if dizzy.</li> <li>• Try to get up and move slowly.</li> </ul>	Contact your health care team if no improvement or if severe
<p><b>Rash; dry, itchy skin</b></p> <ul style="list-style-type: none"> <li>• Stay out of the sun; wear sunblock, a hat and cover exposed skin.</li> <li>• Use daily moisturizer.</li> </ul>	Contact your health care team if no improvement or if severe

Side effects and what to do	When to contact doctor?
<b>Less Common Side Effects, but may be Severe</b>	
<p><b>Allergic reaction</b> (severe rash, itchiness, swollen face, lip or tongue, chest or throat tightness; may occur during or shortly after the drug is given)</p>	Get emergency medical help right away

\*The most updated version and more symptom control information can be found on: <http://www.cancercare.on.ca/druginfo>  
 Prepared with input from the Cancer Care Ontario-Medication Information Sheets Working Group.  
 April 2016

Side effects and what to do	When to contact doctor?
<b>Less Common Side Effects, but may be Severe</b>	
<p><b>Abnormal liver lab tests</b></p> <ul style="list-style-type: none"> <li>Your doctor will monitor these regularly. Call your doctor if you have yellowish skin or eyes, or unusual dark urine.</li> </ul>	Contact your health care team if no improvement or if severe
<p><b>Rupture in stomach or intestine wall</b> (Sudden, severe pain in belly or stomach area)</p>	Get emergency medical help right away
<p><b>Heart problems</b> (irregular heartbeat, chest pain, fainting, swelling, shortness of breath)</p>	Get emergency medical help right away
<p><b>Pancreas problems</b> (increased pain in centre of belly and may extend to back, appetite or weight loss)</p>	Get emergency medical help right away
<p><b>Lung problems</b> (increased cough, breathing problems, chest pain, coughing blood)</p>	Get emergency medical help right away
<p><b>Kidney problems</b> (lower back pain, body swelling, passing little or no urine, or recent unusual weight gain)</p>	Get emergency medical help right away
<p><b>Seizures</b></p>	Get emergency medical help right away
<p><b>Rapid killing of cancer cells when you start treatment may lead to build up of cell waste products</b></p> <ul style="list-style-type: none"> <li>If mild, this may cause gout, with joint pains, but if severe, may cause fevers, kidney failure, confusion and be life-threatening.</li> <li>You <b>MUST</b> take the preventive medicines given by your doctor <b>AND</b></li> <li>Drink plenty of fluids (6-8 glasses per day) and void (urinate) frequently.</li> </ul>	Get emergency medical help right away
<p><b>Blood clot</b> (limb pain or swelling, hardened vein in limb), may occur in lungs (sudden start of coughing, breathing problems, chest pain, coughing blood)</p>	Get emergency medical help right away

For more links on how to manage your symptoms go to [www.cancercare.on.ca/symptoms](http://www.cancercare.on.ca/symptoms).

The information set out in the medication information sheets, regimen information sheets, and symptom management information (for patients) contained in the Drug Formulary (the "Formulary") is intended to be used by health professionals and patients for informational purposes only. The information is not intended to cover all possible uses, directions, precautions, drug interactions or side effects of a certain drug, nor should it be used to indicate that use of a

\*The most updated version and more symptom control information can be found on: <http://www.cancercare.on.ca/druginfo>

Prepared with input from the Cancer Care Ontario-Medication Information Sheets Working Group.

April 2016

*particular drug is safe, appropriate or effective for a given condition.*

*A patient should always consult a healthcare provider if he/she has any questions regarding the information set out in the Formulary. The information in the Formulary is not intended to act as or replace medical advice and should not be relied upon in any such regard. All uses of the Formulary are subject to clinical judgment and actual prescribing patterns may not follow the information provided in the Formulary.*

---

\*The most updated version and more symptom control information can be found on: <http://www.cancercare.on.ca/druginfo>

**Prepared with input from the Cancer Care Ontario-Medication Information Sheets Working Group.**

April 2016