

Medication Information Sheet

dacomitinib (DAK-oh-MI-ti-nib)

This document provides general information about your medication. It does not replace the advice of your health care professional. Always discuss your therapy with your health care professional and refer to the package insert for more details.

Other Name: Vizimpro™

Appearance: tablet in various strengths

What is this medication for?

- For treating a certain type of lung cancer called Non-Small-Cell Lung Cancer (NSCLC)

What should I do before I have this medication?

Tell your health care team if you have or had significant medical condition(s), especially if you have / had:

- frequent diarrhea
- eye problems
- liver, kidney, lung problems (other than lung cancer) or
- any allergies.

Dacomitinib tablets contain a small amount of lactose. If you cannot have lactose, talk to your health care team.

Remember to:

- Tell your health care team about all of the other medications you are taking.
- Keep taking other medications that have been prescribed for you, unless you have been told not to by your health care team.

How will this medication affect sex, pregnancy and breastfeeding?

Talk to your health care team about:

- How this medication may affect your sexual health.
- How this medication may affect your ability to have a baby, if this applies to you.

This medication may harm an unborn baby. Tell your health care team if you or your partner are pregnant, become pregnant during treatment, or are breastfeeding.

- If there is **any** chance of pregnancy happening, you and your partner together must use **2 effective forms of birth control** at the same time for at least **2 months** after your last dose. Talk to your health care team about which birth control options are best for you.
- Do not breastfeed while on this medication and for at least **2 months** after your last dose.

How is this medication given?

- This medication is usually taken once a day by mouth. Talk to your health care team about how and when to take your medication.
- Take the dose at about the same time each day.
- Swallow whole with a glass of water, with or without food.
- Do not crush or chew the tablets.
- If you miss a dose, skip the missed dose. Take your next dose at your usual time. Do not take an extra dose to make up for the missed dose.
- If you vomit (throw up) after taking a dose, do not repeat the dose. Take your next dose at your normal time on the following day.
- If you take too much of your oral anticancer medication by accident, or if you think a child or a pet may have swallowed your medication, you must call the Ontario Poison Control Center right away at: 1-800-268-9017.

To Treat Diarrhea

Dacomitinib can cause diarrhea. Diarrhea is when you have loose bowel movements (watery poo) or you need to go poo (have bowel movements) more often than usual. Diarrhea may start a few days after your treatment.

You may be given a medication called loperamide (Imodium®) to help treat your diarrhea. Take this medication only if you need it.

Keep your loperamide with you all the time. When diarrhea starts, take the loperamide right away and keep taking it until you have no diarrhea for 12 hours.

What else do I need to know while on this medication?

Will this medication interact with other medications or natural health products?

- This medication can interact with other medications, vitamins, foods and natural health products. Interactions can make the treatment not work as well or cause severe side effects.
- Tell your health care team about all of your:
 - ◊ prescription and over-the-counter (non-prescription) medications and all other drugs, such as marijuana (medical or recreational)
 - ◊ natural health products such as vitamins, herbal teas, homeopathic medicines, and other supplements
- Check with your health care team before starting or stopping any of them.

What should I do if I feel unwell, have pain, a headache or a fever?

- **Always** check your temperature to see if you have a fever **before** taking any medications for fever or pain (such as acetaminophen (Tylenol) or ibuprofen (Advil)).
 - ◊ Fever can be a sign of infection that may need treatment right away.
 - ◊ If you take these medications before you check for fever, they may lower your temperature and you may not know you have an infection.

How to check for fever:

Keep a digital (electronic) thermometer at home and take your temperature if you feel hot or unwell (for example, chills, headache, mild pain).

- You have a fever if your temperature taken in your mouth (oral temperature) is:
 - 38.3°C (100.9°F) or higher at any time

OR

 - 38.0°C (100.4°F) or higher for at least one hour.

If you do have a fever:

- **Try to contact your health care team. If you are not able to talk to them for advice, you MUST get emergency medical help right away.**
- Ask your health care team for the [Fever](#) pamphlet for more information.

If you do not have a fever but have mild symptoms such as headache or mild pain:

- Ask your health care team about the right medication for you. **Acetaminophen (Tylenol®)** is a safe choice for most people.
- **Talk to your health care team before you start taking** Ibuprofen (Advil®, Motrin®), naproxen (Aleve®) or ASA (Aspirin®), as they may increase your chance of bleeding or interact with your cancer treatment.
- Talk to your health care team if you already take **low dose aspirin** for a medical condition (such as a heart problem). It may still be safe to take.

What to DO while on this medication:

- DO check with your health care team before getting any vaccinations, surgery, dental work or other medical procedures.
- DO protect your skin from the sun. Wear a long sleeved shirt, long pants and a hat. Apply sunscreen with UVA and UVB protection and an SPF of at least 30. Your skin may be more sensitive to the sun and you could develop a bad sunburn or rash more easily.

What NOT to DO while on this medication:

- DO NOT smoke or drink alcohol while on treatment without talking to your health care team first. Smoking and drinking can make side effects worse and make your treatment not work as well.
- DO NOT drive, operate machinery or do any tasks that need you to be alert if you have any problems with your vision or feel drowsy or dizzy.

How should I safely store this medication?

- Do not throw out any unused medications at home. Bring them to your pharmacy to be thrown away safely.
- Keep this medication in the original packaging at room temperature in a dry place, away from heat and light. Keep out of sight and reach of children and pets.

- **How to safely touch oral anti-cancer medications**

If you are a patient:

- ◊ Wash your hands before and after touching your oral anti-cancer medication.
- ◊ Swallow each pill whole. Do not crush or chew your pills.

If you are a caregiver:

- ◊ Wear nitrile or latex gloves when touching tablets, capsules or liquids.
- ◊ Wash your hands before putting on your gloves and after taking them off, even if your skin did not touch the oral anti-cancer medication.
- ◊ Throw out your gloves after each use. Do not re-use gloves.
- ◊ Do not touch oral anti-cancer medications if you are pregnant or breastfeeding.

- **What to do if oral anti-cancer medication gets on your skin or in your eyes**

If medication gets on your skin:

- ◊ Wash your skin with a lot of soap and water.
- ◊ If your skin gets red or irritated, talk to your health care team.

If medication gets in your eyes:

- ◊ Rinse your eyes with running water right away. Keep water flowing over your open eyes for at least 15 minutes.

What are the side effects of this medication?

The following table lists side effects that you may have when getting dacomitinib. The table is set up to list the most common side effects first and the least common last. It is unlikely that you will have all of the side effects listed and you may have some that are not listed.

Read over the side effect table so that you know what to look for and when to get help. Refer to this table if you experience any side effects while on dacomitinib.

Very Common Side Effects (50 or more out of 100 people)	
Side effects and what to do	When to contact health care team
<p>Diarrhea (may be severe)</p> <p>What to look for?</p> <ul style="list-style-type: none"> Loose, watery, unformed stool (poo) that may happen days to weeks after you get your treatment. <p>What to do?</p> <p>If you have diarrhea:</p> <ul style="list-style-type: none"> Take anti-diarrhea medication if your health care team prescribed it or told you to take it. Do not eat foods or drinks with artificial sweetener (like chewing gum or 'diet' drinks), coffee and alcohol. Eat many small meals and snacks instead of 2 or 3 large meals. Drink at least 6 to 8 cups of liquids each day, unless your health care team has told you to drink more or less. Talk to your health care team if you can't drink 6 to 8 cups of liquids each day when you have diarrhea. You may need to drink special liquids with salt and sugar, called Oral Rehydration Therapy. Talk to your health care team if your diarrhea does not improve after 24 hours of taking diarrhea medication or if you have diarrhea more than 7 times in one day. <p>Ask your health care team for the Diarrhea pamphlet for more information.</p>	<p>Talk to your health care team if no improvement after 24 hours of taking diarrhea medication or if severe (more than 7 times in one day)</p>

Very Common Side Effects (50 or more out of 100 people)	
Side effects and what to do	When to contact health care team
<p>Rash; dry, itchy skin (may be severe)</p> <p>What to look for?</p> <ul style="list-style-type: none"> You may have cracked, rough, flaking or peeling areas of the skin. Your skin may look red and feel warm, like a sunburn. Your skin may itch, burn, sting or feel very tender when touched. <p>What to do?</p> <p>To prevent and treat dry skin:</p> <ul style="list-style-type: none"> Use fragrance-free skin moisturizer. Protect your skin from the sun and the cold. Use sunscreen with UVA and UVB protection and a SPF of at least 30. Avoid perfumed products and lotions that contain alcohol. Drink 6 to 8 cups of non-alcoholic, non-caffeinated liquids each day, unless your health care team has told you to drink more or less. <p>Rash may be severe in some rare cases and cause your skin to blister or peel. If this happens, get emergency medical help right away.</p>	<p>Talk to your health care team if it does not improve or if it is severe</p>
<p>Mouth sores (generally mild)</p> <p>What to look for?</p> <ul style="list-style-type: none"> Round, painful, white or gray sores inside your mouth that can occur on the tongue, lips, gums, or inside your cheeks. In more severe cases they may make it hard to swallow, eat or brush your teeth. They may last for 3 days or longer. <p>What to do?</p> <p>To help prevent mouth sores:</p> <ul style="list-style-type: none"> Take care of your mouth by gently brushing and flossing regularly. Rinse your mouth often with a homemade mouthwash. To make a homemade mouthwash, mix 1 teaspoonful of baking soda and 1 teaspoonful of salt in 4 cups (1L) of water. 	<p>Talk to your health care team as soon as you notice mouth or lip sores or if it hurts to eat, drink or swallow</p>

Very Common Side Effects (50 or more out of 100 people)	
Side effects and what to do	When to contact health care team
<ul style="list-style-type: none"> Do not use store-bought mouthwashes, especially those with alcohol, because they may irritate your mouth. <p>If you have mouth sores:</p> <ul style="list-style-type: none"> Avoid hot, spicy, acidic, hard or crunchy foods. Your doctor may prescribe a special mouthwash to relieve mouth sores and prevent infection. Talk to your health care team as soon as you notice mouth or lip sores or if it hurts to eat, drink or swallow. <p>Ask your health care team for the Oral Care (Mouth Care) pamphlet for more information.</p>	
<p>Nail changes (may be severe)</p> <p>What to look for?</p> <ul style="list-style-type: none"> You may have changes in nail colour, pain or tenderness, swelling of cuticles, or loosening of nails. Nails will slowly return to normal after treatment ends. <p>What to do?</p> <ul style="list-style-type: none"> Moisturize your nails and cuticles. Do not use nail polish and fake fingernails until your nails have gone back to normal. Wear gloves when doing house chores or gardening. 	Talk to your health care team if it does not improve or if it is severe

Common Side Effects (25 to 49 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>Too much or too little salt in your body</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Muscle spasms, cramping, weakness, twitching, or convulsions. • Irregular heartbeat, confusion or blood pressure changes. <p>What to do?</p> <p>Get emergency medical help right away for severe symptoms.</p>	<p>Get emergency medical help right away for severe symptoms</p>
<p>Low appetite</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Loss of interest in food or not feeling hungry. • Weight loss. <p>What to do?</p> <ul style="list-style-type: none"> • Try to eat your favourite foods. • Eat small meals throughout the day. • You may need to take meal supplements to help keep your weight up. • Talk to your health care team if you have no appetite. <p>Ask your health care team for the Loss of Appetite pamphlet for more information.</p>	<p>Talk to your health care team if it does not improve or if it is severe</p>

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>Liver problems (may be severe)</p> <p>Your health care team may check your liver function with a blood test. The liver changes do not usually cause any symptoms.</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Rarely, you may develop yellowish skin or eyes, unusually dark pee or pain on the right side of your belly. This may be severe. <p>What to do?</p> <p>If you have any symptoms of liver problems, get emergency medical help right away.</p>	<p>Get emergency medical help right away</p>
<p>Eye problems (may be severe)</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Your eyes may feel dry, irritated, or painful. • They may look red and have a lot of tears. • They may feel sensitive to light and your vision may be blurry. <p>What to do?</p> <ul style="list-style-type: none"> • Avoid wearing contact lenses. • Wear sunglasses with UV protection. • Use protective eyewear (goggles or helmet with face mask) when playing sports, mowing the lawn or doing anything that may get particles or fumes in your eyes. • You may try artificial tears (eye drops) or ointment. 	<p>Talk to your health care team as soon as possible</p>
<p>Hair thinning or loss</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Your hair may begin to become thin or fall out during or after treatment. • In most cases, your hair will grow back after treatment, but the texture or colour may change. • In very rare cases, hair loss may be permanent. 	<p>Talk to your health care team if this bothers you</p>

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>What to do?</p> <ul style="list-style-type: none"> • Use a gentle soft brush. • Do not use hair sprays, bleaches, dyes and perms. 	
<p>Cough and feeling short of breath</p> <p>What to look for?</p> <ul style="list-style-type: none"> • You may have a cough and feel short of breath. • Symptoms that commonly occur with a cough are: <ul style="list-style-type: none"> ◦ Wheezing or a whistling breathing ◦ Runny nose ◦ Sore throat ◦ Heartburn ◦ Weight loss ◦ Fever and chills • Rarely this may be severe with chest pain, trouble breathing or coughing up blood. <p>What to do?</p> <ul style="list-style-type: none"> • Check your temperature to see if you have a fever. Read the above section "What should I do if I feel unwell, have pain, a headache or a fever?" • If you have a fever, try to talk to your health care team. If you are not able to talk to them for advice, you MUST get emergency medical help right away. • If you have a severe cough with chest pain, trouble breathing or you are coughing up blood, get medical help right away. 	<p>Talk to your health care team. If you are not able to talk to your health care team for advice, and you have a fever or severe symptoms, you MUST get emergency medical help right away</p>
<p>Nausea and vomiting (generally mild)</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Nausea is feeling like you need to throw up. You may also feel light-headed. • You may feel nausea within hours to days after your treatment. 	<p>Talk to your health care team if nausea lasts more than 48 hours or vomiting lasts more than 24 hours or if severe</p>

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>What to do?</p> <p>To help prevent nausea:</p> <ul style="list-style-type: none"> • It is easier to prevent nausea than to treat it once it happens. • Drink clear liquids and have small meals. Get fresh air and rest. • Do not eat spicy, fried foods or foods with a strong smell. • Limit caffeine (like coffee, tea) and avoid alcohol. <p>If you have nausea or vomiting:</p> <ul style="list-style-type: none"> • Take your rescue (as-needed) anti-nausea medication(s) as prescribed. • Ask your health care team for the Nausea & Vomiting pamphlet for more information. • Talk to your health care team if: <ul style="list-style-type: none"> ◦ nausea lasts more than 48 hours ◦ vomiting lasts more than 24 hours or if it is severe 	
<p>Rash on your hands and feet (hand-foot syndrome)</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Tingling or swelling of the skin on the palms of your hands and the bottoms of your feet. This can become painful, red and numb. • In worse cases your skin may start to peel and you can get blisters or sores. • This may happen days or weeks after you start treatment. <p>What to do?</p> <p>To help prevent Hand-foot syndrome:</p> <ul style="list-style-type: none"> • Do not do activities that cause rubbing or pressure on your skin, like heavy-duty washing, gripping tools, typing, playing musical instruments, and driving. • Moisturize your hands and feet often, especially in the skin folds. • Wear loose, comfortable footwear and clothes. • Rest and try to keep off your feet. • Do not let your hands and feet get too hot. <p>Also your health care team for the Hand-foot syndrome pamphlet for more information.</p>	<p>Talk to your health care team if it does not improve or if it is severe</p>

The most updated information sheet version can be found on <https://www.cancercareontario.ca/drugs>
 Additional symptom management information is available from <https://www.cancercareontario.ca/symptoms>
 Created by the CCO Drug Formulary team, with input from the CCO Patient Education team and Patient & Family Advisors.
 November 2020

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>Fatigue</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Feeling of tiredness or low energy that lasts a long time and does not go away with rest or sleep. <p>What to do?</p> <ul style="list-style-type: none"> • Be active. Aim to get 30 minutes of moderate exercise (you are able to talk comfortably while exercising) on most days. • Check with your health care team before starting any new exercise. • Pace yourself, do not rush. Put off less important activities. Rest when you need to. • Ask family or friends to help you with things like housework, shopping, and child or pet care. • Eat well and drink at least 6 to 8 glasses of water or other liquids every day (unless your health care team has told you to drink more or less). • Avoid driving or using machinery if you are feeling tired. <p>Ask your health care team for the Fatigue pamphlet for more information.</p>	<p>Talk to your health care team if it does not improve or if it is severe</p>
<p>Mild joint, muscle pain or cramps</p> <p>What to look for?</p> <ul style="list-style-type: none"> • New pain in your muscles or joints, muscle cramps, or feeling achy. <p>What to do?</p> <ul style="list-style-type: none"> • Take pain medication (acetaminophen or opioids such as codeine, morphine, hydromorphone, oxycodone) as prescribed. • Read the above section: "What should I do if I feel unwell, have pain, a headache or a fever?" before taking acetaminophen (Tylenol®), ibuprofen (Advil®, Motrin®), naproxen (Aleve®) or Aspirin. These medications may hide an infection that needs treatment or they may increase your risk of bleeding. • Rest often and try light exercise (such as walking) as it may help. <p>Ask your health care team for the Pain pamphlet for more information.</p>	<p>Talk to your health care team if it does not improve or if it is severe</p>

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>Trouble Sleeping</p> <p>Your medications may cause trouble sleeping. It may get better once your body gets used to the medication or when your treatment ends.</p> <p>What to look for?</p> <ul style="list-style-type: none"> • You may find it hard to fall asleep or stay asleep. • How well you sleep may change over your treatment. For example, you may have several nights of poor sleep followed by a night of better sleep. • You may wake up too early or not feel well-rested after a night's sleep. • You may feel tired or sleepy during the day. <p>What to do?</p> <p>Talk to your health care team if it does not improve or if it is severe.</p>	<p>Talk to your health care team if it does not improve or if it is severe</p>

Who do I contact if I have questions or need help?

My cancer health care provider is: _____

During the day I should contact: _____

Evenings, weekends and holidays: _____

Other Notes:

November 2020 New info sheet

For more links on how to manage your symptoms go to www.cancercareontario.ca/symptoms.

The information set out in the medication information sheets, regimen information sheets, and symptom management information (for patients) contained in the Drug Formulary (the "Formulary") is intended to be used by health professionals and patients for informational purposes only. The information is not intended to cover all possible uses, directions, precautions, drug interactions or side effects of a certain drug, nor should it be used to indicate that use of a particular drug is safe, appropriate or effective for a given condition.

A patient should always consult a healthcare provider if he/she has any questions regarding the information set out in the Formulary. The information in the Formulary is not intended to act as or replace medical advice and should not be relied upon in any such regard. All uses of the Formulary are subject to clinical judgment and actual prescribing patterns may not follow the information provided in the Formulary.