

Medication Information Sheet

gemcitabine (jem-SITE-a-been)

This document provides general information about your medication. It does not replace the advice of your health care professional. Always discuss your therapy with your health care professional and refer to the package insert for more details.

Other Name: Gemzar®

Appearance: Clear, colourless solution mixed into larger bags of fluids

What is this medication for?

- For treating certain types of cancers such as bladder, pancreas, lung, breast and other types of cancer.

What should I do before I have this medication?

- Tell your health care team if you have or had significant medical condition(s), especially if you have or had:
 - liver, lung, kidney or heart problems
 - Issues with your bone marrow or
 - any allergies.
- Tell your health care team if you have had or will be getting radiation therapy.

Remember to:

- Tell your health care team about all of the other medications you are taking.
- Keep taking other medications that have been prescribed for you, unless you have been told not to by your health care team.

How will this medication affect sex, pregnancy and breastfeeding?**Talk to your health care team about:**

- How this medication may affect your sexual health.
- How this medication may affect your ability to have a baby, if this applies to you.

This medication may harm an unborn baby. Tell your health care team if you or your partner are pregnant, become pregnant during treatment, or are breastfeeding.

- If there is **any** chance of pregnancy happening, you and your partner together must use **2 effective forms of birth control** at the same time until **6 months** after your last dose. Talk to your health care team about which birth control options are best for you.
- Do not use hormonal birth control (such as birth control pills), unless your health care team told you that they are safe. Talk to your health care team about the safest birth control for you.
- Do not breastfeed while on this medication.

How is this medication given?

- This drug is given through an IV (injected into a vein) . Talk to your health care team about your treatment schedule.
- If you missed your treatment appointment, talk to your health care team to find out what to do.

To Prevent or Treat Nausea and Vomiting

You may be given medications to help prevent nausea (feeling like throwing up) and vomiting (throwing up) before they start.

- These are called anti-nausea medications and include medications such as dexamethasone or others.

What else do I need to know while on this medication?

- **Will this medication interact with other medications or natural health products?**
 - Although this medication is unlikely to interact with other medications, vitamins, foods and natural health products, tell your health care team about all of your:
 - prescription and over-the-counter (non-prescription) medications and all other drugs, such as marijuana (medical or recreational)
 - natural health products such as vitamins, herbal teas, homeopathic medicines, and other supplements
 - Check with your health care team before starting or stopping any of them.
- If you are taking a blood thinner (such as warfarin), your health care team may need extra blood tests and may change your dose.

- **What should I do if I feel unwell, have pain, a headache or a fever?**
 - **Always** check your temperature to see if you have a fever **before** taking any medications for fever or pain (such as acetaminophen (Tylenol) or ibuprofen (Advil)).
 - Fever can be a sign of infection that may need treatment right away.
 - If you take these medications before you check for fever, they may lower your temperature and you may not know you have an infection.

How to check for fever:

Keep a digital (electronic) thermometer at home and take your temperature if you feel hot or unwell (for example, chills, headache, mild pain).

- You have a fever if your temperature taken in your mouth (oral temperature) is:
 - 38.3°C (100.9°F) or higher at any time

OR

 - 38.0°C (100.4°F) or higher for at least one hour.

If you do have a fever:

- **Try to contact your health care team. If you are not able to talk to them for advice, you MUST get emergency medical help right away.**
- Ask your health care team for the [Fever](#) pamphlet for more information.

If you do not have a fever but have mild symptoms such as headache or mild pain:

- Ask your health care team about the right medication for you. **Acetaminophen (Tylenol®)** is a safe choice for most people.
- **Talk to your health care team before you start taking** Ibuprofen (Advil®, Motrin®), naproxen (Aleve®) or ASA (Aspirin®), as they may increase your chance of bleeding or interact with your cancer treatment.
- Talk to your health care team if you already take **low dose aspirin** for a medical condition (such as a heart problem). It may still be safe to take.

- **What to DO while on this medication:**

- DO check with your health care team before getting any vaccinations, surgery, dental work or other medical procedures.

- DO consider asking someone to drive you to and from the hospital on your treatment days. You may feel weak or dizzy after your treatment.
- **What NOT to DO on this medication:**
- DO NOT smoke or drink alcohol while on treatment without talking to your health care team first. Smoking and drinking can make side effects worse and make your treatment not work as well.

NOTE:

- Gemcitabine makes your body more sensitive to radiation therapy. If you get radiation treatment at the same time as gemcitabine, you may get worse side effects from radiation.

What are the side effects of this medication?

The following table lists side effects that you may have when getting gemcitabine. The table is set up to list the most common side effects first and the least common last. It is unlikely that you will have all of the side effects listed and you may have some that are not listed.

Read over the side effect table so that you know what to look for and when to get help. Refer to this table if you experience any side effects while on gemcitabine.

Very Common Side Effects (50 or more out of 100 people)	
Side effects and what to do	When to contact health care team
<p>Low neutrophils (white blood cells) in the blood (neutropenia)</p> <p>(May be severe)</p> <p>When neutrophils are low, you are at risk of getting an infection more easily. Ask your health care team for the Neutropenia (Low white blood cell count) pamphlet for more information.</p> <p>What to look for?</p> <ul style="list-style-type: none"> • If you feel hot or unwell (for example if you have chills or a new cough), you must check your temperature to see if you have a fever. • Do not take medications that treat a fever before you take your temperature (for example, Tylenol®, acetaminophen, Advil® or ibuprofen). • Do not eat or drink anything hot or cold right before taking your temperature. <p>You have a fever if your temperature taken in your mouth (oral temperature) is:</p> <ul style="list-style-type: none"> • 38.3°C (100.9°F) or higher at any time OR • 38.0°C (100.4°F) or higher for at least one hour. <p>What to do?</p> <p>If your health care team has told you that you have low neutrophils:</p> <ul style="list-style-type: none"> • Wash your hands often to prevent infection. • Check with your health care team before getting any vaccines, surgeries, medical procedures or visiting your dentist. • Keep a digital thermometer at home so you can easily check for a fever. <p>If you have a fever:</p> <p>If you have a fever, try to contact your health care team. If you are unable to talk to the team for advice, you must get emergency medical help right away.</p>	<p>If you have a fever, try to contact your health care team. If you are unable to talk to the team for advice, you MUST get emergency medical help right away.</p>

Very Common Side Effects (50 or more out of 100 people)	
Side effects and what to do	When to contact health care team
<p>Low platelets in the blood</p> <p>(May be severe)</p> <p>When your platelets are low, you are at risk for bleeding and bruising. Ask your health care team for the Low Platelet Count pamphlet for more information.</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Watch for signs of bleeding: <ul style="list-style-type: none"> ◦ bleeding from your gums ◦ unusual or heavy nosebleeds ◦ bruising easily or more than normal ◦ black coloured stools (poo) or blood in your stools (poo) ◦ coughing up red or brown coloured mucus ◦ dizziness, constant headache or changes in your vision ◦ heavy vaginal bleeding ◦ red or pink coloured urine (pee) <p>What to do?</p> <p>If your health care team has told you that you have low platelets:</p> <ul style="list-style-type: none"> • Tell your pharmacist that your platelet count may be low before taking any prescriptions or over-the-counter medication. • Check with your healthcare team before you go to the dentist. • Take care of your mouth and use a soft toothbrush. • Try to prevent cuts and bruises. • Ask your health care team what activities are safe for you. • Your treatment may have to be delayed if you have low platelets. Your health care team may recommend a blood transfusion. <p>If you have signs of bleeding:</p> <ul style="list-style-type: none"> • If you have a small bleed, clean the area with soap and water or a saline (saltwater) rinse. Apply pressure for at least 10 minutes. <p>If you have bleeding that does not stop or is severe (very heavy), you must get emergency medical help right away.</p>	<p>Talk to your health care team if you have any signs of bleeding. If you have bleeding that doesn't stop or is severe (very heavy), you MUST get emergency help right away.</p>

Very Common Side Effects (50 or more out of 100 people)	
Side effects and what to do	When to contact health care team
<p>Liver problems</p> <p>Your health care team may check your liver function with a blood test. The liver changes do not usually cause any symptoms.</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Rarely, you may develop yellowish skin or eyes, unusually dark pee or pain on the right side of your belly. This may be severe. <p>What to do?</p> <p>If you have any symptoms of liver problems, get emergency medical help right away.</p>	<p>Get emergency medical help right away</p>
<p>Nausea and vomiting</p> <p>(Generally mild)</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Nausea is feeling like you need to throw up. You may also feel light-headed. • You may feel nausea within hours to days after your treatment. <p>What to do?</p> <p>To help prevent nausea:</p> <ul style="list-style-type: none"> • It is easier to prevent nausea than to treat it once it happens. • Take your anti-nausea medication(s) as prescribed, even if you do not feel like throwing up. • Drink clear liquids and have small meals. Get fresh air and rest. • Do not eat spicy, fried foods or foods with a strong smell. • Limit caffeine (like coffee, tea) and avoid alcohol. <p>If you have nausea or vomiting:</p> <ul style="list-style-type: none"> • Take your rescue (as-needed) anti-nausea medication(s) as prescribed. • Ask your health care team for the Nausea & Vomiting pamphlet for more information. • Talk to your health care team if: <ul style="list-style-type: none"> ◦ nausea lasts more than 48 hours ◦ vomiting lasts more than 24 hours or if it is severe 	<p>Talk to your health care team if nausea lasts more than 48 hours or vomiting lasts more than 24 hours or if severe</p>

Common Side Effects (25 to 49 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>Headache, Flu-like symptoms</p> <p>You may feel like you have the flu for around 2 days after your IV treatment. These flu-like symptoms may not be signs of an infection.</p> <p>What to look for?</p> <ul style="list-style-type: none"> You may have chills, headache and muscle pain. You may feel tired and have a poor appetite. Symptoms may happen at any time after you receive your treatment and usually go away as your body gets used to the medication. <p>What to do?</p> <ul style="list-style-type: none"> Check your temperature to see if you have a fever. Read the above section "What should I do if I feel unwell, have pain, a headache or a fever?" on page 3. If you do have a fever, try to speak to your health care team. If you are unable to talk to them for advice, you MUST get emergency medical help right away. 	<p>Talk to your health care team if it does not improve or if it is severe</p>
<p>Proteins in Urine</p> <p>Your health care team may do urine tests to check for proteins in your pee.</p> <p>What to look for?</p> <ul style="list-style-type: none"> Swelling in your face, legs, or belly. Recent weight gain that is not normal for you. Foamy, frothy, or bubbly-looking pee. <p>What to do?</p> <p>Talk to your health care team if it does not improve or if it is severe.</p>	<p>Talk to your health care team if it does not improve or if it is severe</p>
<p>Rash; dry, itchy skin</p> <p>(Rarely severe)</p> <p>What to look for?</p> <ul style="list-style-type: none"> You may have cracked, rough, flaking or peeling areas of the skin. 	<p>Talk to your health care team if it does not improve or if it is severe</p>

Common Side Effects (25 to 49 out of 100 people)	
Side effects and what to do	When to contact health care team
<ul style="list-style-type: none"> Your skin may look red and feel warm, like a sunburn. Your skin may itch, burn, sting or feel very tender when touched. <p>What to do?</p> <p>To prevent and treat dry skin:</p> <ul style="list-style-type: none"> Use fragrance-free skin moisturizer. Protect your skin from the sun and the cold. Use sunscreen with UVA and UVB protection and a SPF of at least 30. Avoid perfumed products and lotions that contain alcohol. Drink 6 to 8 cups of non-alcoholic, non-caffeinated liquids each day, unless your health care team has told you to drink more or less. <p>Rash may be severe in some rare cases and cause your skin to blister or peel. If this happens, get emergency medical help right away.</p>	

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>Mild swelling</p> <p>What to look for?</p> <ul style="list-style-type: none"> You may have mild swelling or puffiness in your arms and/or legs. Rarely, this may be severe. <p>What to do?</p> <p>To help prevent swelling:</p> <ul style="list-style-type: none"> Eat a low-salt diet. <p>If you have swelling:</p> <ul style="list-style-type: none"> Wear loose-fitting clothing. For swollen legs or feet, keep your feet up when sitting. 	<p>Talk to your health care team if it does not improve or if it is severe</p>

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>Mild joint, muscle pain or cramps</p> <p>What to look for?</p> <ul style="list-style-type: none"> • New pain in your muscles or joints, muscle cramps, or feeling achy. <p>What to do?</p> <ul style="list-style-type: none"> • Take pain medication (acetaminophen or opioids such as codeine, morphine, hydromorphone, oxycodone) as prescribed. • Read the above section: "What should I do if I feel unwell, have pain, a headache or a fever?" on page 3 before taking acetaminophen (Tylenol®), ibuprofen (Advil®, Motrin®), naproxen (Aleve®) or Aspirin. These medications may hide an infection that needs treatment or they may increase your risk of bleeding. • Rest often and try light exercise (such as walking) as it may help. <p>Ask your health care team for the Pain pamphlet for more information.</p>	<p>Talk to your health care team if it does not improve or if it is severe</p>
<p>Hair thinning or loss</p> <p>(Generally mild)</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Your hair may begin to become thin or fall out during or after treatment. • In most cases, your hair will grow back after treatment, but the texture or colour may change. • In very rare cases, hair loss may be permanent. <p>What to do?</p> <ul style="list-style-type: none"> • Use a gentle soft brush. • Do not use hair sprays, bleaches, dyes and perms. 	<p>Talk to your health care team if this bothers you</p>
<p>Diarrhea</p> <p>What to look for?</p> <ul style="list-style-type: none"> • Loose, watery, unformed stool (poo) that may happen days to weeks after you get your treatment. 	<p>Talk to your health care team if no improvement after 24 hours of taking diarrhea medication or if severe (more than 7 times in one day)</p>

Less Common Side Effects (10 to 24 out of 100 people)	
Side effects and what to do	When to contact health care team
<p>What to do?</p> <p>If you have diarrhea:</p> <ul style="list-style-type: none"> • Take anti-diarrhea medication if your health care team prescribed it or told you to take it. • Do not eat foods or drinks with artificial sweetener (like chewing gum or 'diet' drinks), coffee and alcohol. • Eat many small meals and snacks instead of 2 or 3 large meals. • Drink at least 6 to 8 cups of liquids each day, unless your health care team has told you to drink more or less. • Talk to your health care team if you can't drink 6 to 8 cups of liquids each day when you have diarrhea. You may need to drink special liquids with salt and sugar, called Oral Rehydration Therapy. • Talk to your health care team if your diarrhea does not improve after 24 hours of taking diarrhea medication or if you have diarrhea more than 7 times in one day. <p>Ask your health care team for the Diarrhea pamphlet for more information.</p>	

- Sudden onset of signs and symptoms of kidney failure such as leg/body swelling or passing little or no urine
- Sudden changes to vision, speech, or the use of your limb(s)
- Chest pain, shortness of breath or pain in chest, belly or limb
- Increased cough or coughing up blood
- Dizziness or passing out, unexpected weight gain, throwing up blood or throw up that looks like coffee grounds or black, tarry, or bloody stools.
- Irregular heartbeat or chest pain
- Severe swollen face, lip or tongue, chest or throat tightness; may occur during or shortly after the drug is given
- Red or purple patches on the skin, rash/patches that do not turn white when pressed
- Severe headache, loss of consciousness, seizures, confusion, vision loss

Who do I contact if I have questions or need help?

My cancer health care provider is: _____

During the day I should contact: _____

Evenings, weekends and holidays: _____

Other Notes:

April 2020 Updated side effects and changed to new format.

For more links on how to manage your symptoms go to www.cancercareontario.ca/symptoms.

The information set out in the medication information sheets, regimen information sheets, and symptom management information (for patients) contained in the Drug Formulary (the "Formulary") is intended to be used by health professionals and patients for informational purposes only. The information is not intended to cover all possible uses, directions, precautions, drug interactions or side effects of a certain drug, nor should it be used to indicate that use of a particular drug is safe, appropriate or effective for a given condition.

A patient should always consult a healthcare provider if he/she has any questions regarding the information set out in the Formulary. The information in the Formulary is not intended to act as or replace medical advice and should not be relied upon in any such regard. All uses of the Formulary are subject to clinical judgment and actual prescribing patterns may not follow the information provided in the Formulary.