

## Medication Information Sheet

**pomalidomide** ( Pom-a-LID-oh-mide)

*This document provides general information about your medication. It does not replace the advice of your health care professional. Always discuss your therapy with your health care professional and refer to the package insert for more details.*

**Other Name:** Pomalyst®

**Appearance:** capsule in various strengths and colours

**What is this medication for?**

- For treating a type of blood cancer called multiple myeloma

**What should I do before I have this medication?**

- You must register with the RevAid® program before starting pomalidomide. Your doctor and pharmacy must also be registered with this program for you to receive pomalidomide.
- Tell your doctor and pharmacist if you have or had significant medical condition(s), especially if you have or had:
  - ◊ blood clots or severe bleeding,
  - ◊ severe infections, especially hepatitis B infection,
  - ◊ liver, kidney, heart or nerve problems,
  - ◊ a reaction to thalidomide or lenalidomide, or
  - ◊ any allergies.
- Tell your doctor and pharmacist if you take hormonal drugs, such as birth control pills, or if you smoke cigarettes.

**Remember to:**

- Tell your health care team about all of the other medications you are taking.
- Keep taking other medications that have been prescribed for you, unless you have been told not to by your health care team.

## How will this medication affect sex, pregnancy and breastfeeding?

### Talk to your health care team about:

- How this medication may affect your sexual health.
- How this medication may affect your ability to have a baby, if this applies to you.

This medication may harm an unborn baby. Tell your health care team if you or your partner are pregnant, become pregnant during treatment, or are breastfeeding.

- If there is **any** chance of pregnancy happening, you and your partner together must:
  - ▶ **Use 2 effective forms of birth control** at the same time until **4 weeks** after your last dose **AND have regular pregnancy tests** (see the RevAid® program for full details of requirements).
- Do not use hormonal birth control (such as birth control pills), as these can increase your risk of blood clots. Male patients should use a condom (even if they have had a vasectomy) during sex with females who may become pregnant. Talk to your health care team about which birth control options are best for you.
- Do not breastfeed while on this medication.

## How is this medication given?

- This medication is usually taken once a day by mouth on certain days within a treatment cycle. Talk to your health care team about how and when to take your pomalidomide.
- Pomalidomide is usually given together with other medications. Talk to your health care team about your treatment schedule.
- Swallow pomalidomide capsules whole with a glass of water, with or without food. Do not crush or open capsules.
- If you vomit (throw up) after taking your medication, talk to your health care team about what to do.
- If you miss a dose, take it if it is within 12 hours from the missed dose, otherwise skip this and take your next dose as scheduled. Do not double the dose to make up for the forgotten one.
- If you have kidney problems and are receiving hemodialysis, take your dose after hemodialysis, on hemodialysis days.
- If you take too much of your oral anticancer medication by accident, or if you think a child or a pet may have swallowed your medication, you must call the Ontario Poison Control Center right away.

### To Prevent Blood Clots

This medication may increase the risk of blood clots in some people. Ask your health care team if you are at risk for blood clots.

If you are at risk for blood clots, you may be given blood thinners such as low dose Aspirin®, low molecular weight heparins (such as dalteparin (Fragmin®) or enoxaparin (Lovenox®)), or warfarin (Coumadin®).

### To Prevent Tumor Lysis Syndrome (TLS)

TLS can happen when a large number of cancer cells die quickly and your body cannot get rid of them fast enough. TLS can make you very sick. Ask your health care team if you are at risk for TLS.

If you are at risk for TLS, you may be given medications before your pomalidomide treatment to help prevent it.

These are called anti-uricemics (such as allopurinol), or others.

### To Prevent Hepatitis B Flare Ups

If you have ever been infected with hepatitis B, there is a risk that this treatment can cause it to flare up (come back). Tell your health care team if you have had hepatitis B. You may need to take medication to prevent a hepatitis B flare-up.

## What else do I need to know while on this medication?

- **Will this medication interact with other medications or natural health products?**
  - ◊ This medication can interact with other medications, vitamins, foods and natural health products. Interactions can make the treatment not work as well or cause severe side effects.
  - ◊ Tell your health care team about all of your:
    - prescription and over-the-counter (non-prescription) medications and all other drugs, such as marijuana (medical or recreational)
    - natural health products such as vitamins, herbal teas, homeopathic medicines, and other supplements
  - ◊ Check with your health care team before starting or stopping any of them.

- **What should I do if I feel unwell, have pain, a headache or a fever?**

- **Always** check your temperature to see if you have a fever **before** taking any medications for fever or pain (such as acetaminophen (Tylenol) or ibuprofen (Advil)).
  - Fever can be a sign of infection that may need treatment right away.
  - If you take these medications before you check for fever, they may lower your temperature and you may not know you have an infection.

**How to check for fever:**

Keep a digital (electronic) thermometer at home and take your temperature if you feel hot or unwell (for example, chills, headache, mild pain).

- You have a fever if your temperature taken in your mouth (oral temperature) is:
    - 38.3°C (100.9°F) or higher at any time
- OR
- 38.0°C (100.4°F) or higher for at least one hour.

**If you do have a fever:**

- **Try to contact your health care team. If you are not able to talk to them for advice, you MUST get emergency medical help right away.**
- Ask your health care team for the [Fever](#) pamphlet for more information.

**If you do not have a fever** but have mild symptoms such as headache or mild pain:

- Ask your health care team about the right medication for you. **Acetaminophen (Tylenol®)** is a safe choice for most people.
- **Talk to your health care team before you start taking** Ibuprofen (Advil®, Motrin®), naproxen (Aleve®) or ASA (Aspirin®), as they may increase your chance of bleeding or interact with your cancer treatment.
- Talk to your health care team if you already take **low dose aspirin** for a medical condition (such as a heart problem). It may still be safe to take.

**What to DO while on this medication:**

- DO check with your health care team before getting any vaccinations, surgery, dental work or other medical procedures.
- DO talk to your health care team about your risk of getting other cancers and heart problems after this treatment.
- DO tell your health care team about any serious infections that you have now or have had in the past.

**What NOT to DO while on this medication:**

- DO NOT smoke or drink alcohol while on treatment without talking to your health care team first. Smoking and drinking can make side effects worse and make your treatment not work as well.

## How should I safely store this medication?

- Do not throw out any unused medications at home. Bring them to your pharmacy to be thrown away safely.
- Keep this medication in the original packaging at room temperature in a dry place, away from heat and light. Keep out of sight and reach of children and pets.
- **How to safely touch oral anti-cancer medications**

**If you are a patient:**

- ◊ Wash your hands before and after touching your oral anti-cancer medication.
- ◊ Swallow each pill whole. Do not crush or chew your pills.

**If you are a caregiver:**

- ◊ Wear nitrile or latex gloves when touching tablets, capsules or liquids.
- ◊ Wash your hands before putting on your gloves and after taking them off, even if your skin did not touch the oral anti-cancer medication.
- ◊ Throw out your gloves after each use. Do not re-use gloves.
- ◊ Do not touch oral anti-cancer medications if you are pregnant or breastfeeding.

- **What to do if oral anti-cancer medication gets on your skin or in your eyes**

**If medication gets on your skin:**

- Wash your skin with a lot of soap and water. .
- If your skin gets red or irritated, talk to your health care team.

**If medication gets in your eyes:**

- Rinse your eyes with running water right away. Keep water flowing over your open eyes for at least 15 minutes.

## What are the side effects of this medication?

The following table lists side effects that you may have when getting pomalidomide. The table is set up to list the most common side effects first and the least common last. It is unlikely that you will have all of the side effects listed and you may have some that are not listed.

Read over the side effect table so that you know what to look for and when to get help. Refer to this table if you experience any side effects while on pomalidomide.

Common Side Effects (25 to 49 out of 100 people)	
Side effects and what to do	When to contact health care team
<p><b>Low neutrophils (white blood cells) in the blood (neutropenia) (may be severe)</b></p> <p>When neutrophils are low, you are at risk of getting an infection more easily. Ask your health care team for the <a href="#">Neutropenia (Low Neutrophil Count)</a> pamphlet for more information.</p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• If you feel hot or unwell (for example if you have chills or a new cough), you <b>must</b> check your temperature to see if you have a fever.</li> <li>• Do not take medications that treat a fever before you take your temperature (for example, Tylenol®, acetaminophen, Advil® or ibuprofen).</li> <li>• Do not eat or drink anything hot or cold right before taking your temperature.</li> </ul>	<p>If you have a fever, try to contact your health care team. If you are unable to talk to the team for advice, you <b>MUST</b> get emergency medical help right away.</p>

<b>Common Side Effects (25 to 49 out of 100 people)</b>	
<p>You have a fever if your temperature taken in your mouth (oral temperature) is:</p> <ul style="list-style-type: none"> <li>• <b>38.3°C (100.9°F) or higher at any time</b></li> </ul> <p><b>OR</b></p> <ul style="list-style-type: none"> <li>• <b>38.0°C (100.4°F) or higher for at least one hour.</b></li> </ul> <p><b>What to do?</b></p> <p><b>If your health care team has told you that you have low neutrophils:</b></p> <ul style="list-style-type: none"> <li>• Wash your hands often to prevent infection.</li> <li>• Check with your health care team before getting any vaccines, surgeries, medical procedures or visiting your dentist.</li> <li>• Keep a digital thermometer at home so you can easily check for a fever.</li> </ul> <p><b>If you have a fever:</b></p> <p><b>If you have a fever, try to contact your health care team. If you are unable to talk to the team for advice, you must get emergency medical help right away.</b></p>	
<p><b>Low platelets in the blood (may be severe)</b></p> <p>When your platelets are low, you are at risk for bleeding and bruising. Ask your health care team for the <a href="#">Low Platelet Count</a> pamphlet for more information.</p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• Watch for signs of bleeding: <ul style="list-style-type: none"> <li>◦ bleeding from your gums</li> <li>◦ unusual or heavy nosebleeds</li> <li>◦ bruising easily or more than normal</li> <li>◦ black coloured stools (poo) or blood in your stools (poo)</li> <li>◦ coughing up red or brown coloured mucus</li> <li>◦ dizziness, constant headache or changes in your vision</li> <li>◦ heavy vaginal bleeding</li> <li>◦ red or pink coloured urine (pee)</li> </ul> </li> </ul>	<p>Talk to your health care team if you have any signs of bleeding. If you have bleeding that doesn't stop or is severe (very heavy), you <b>MUST</b> get emergency medical help right away.</p>

<b>Common Side Effects (25 to 49 out of 100 people)</b>	
<p><b>What to do?</b></p> <p><b>If your health care team has told you that you have low platelets:</b></p> <ul style="list-style-type: none"> <li>• Tell your pharmacist that your platelet count may be low before taking any prescriptions or over-the-counter medication.</li> <li>• Check with your healthcare team before you go to the dentist.</li> <li>• Take care of your mouth and use a soft toothbrush.</li> <li>• Try to prevent cuts and bruises.</li> <li>• Ask your health care team what activities are safe for you.</li> <li>• Your treatment may have to be delayed if you have low platelets. Your health care team may recommend a blood transfusion.</li> </ul> <p><b>If you have signs of bleeding:</b></p> <ul style="list-style-type: none"> <li>• If you have a small bleed, clean the area with soap and water or a saline (saltwater) rinse. Apply pressure for at least 10 minutes.</li> </ul> <p><b>If you have bleeding that does not stop or is severe (very heavy), you must get emergency medical help right away.</b></p>	
<p><b>Fatigue</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• Feeling of tiredness or low energy that lasts a long time and does not go away with rest or sleep.</li> </ul> <p><b>What to do?</b></p> <ul style="list-style-type: none"> <li>• Be active. Aim to get 30 minutes of moderate exercise (you are able to talk comfortably while exercising) on most days.</li> <li>• Check with your health care team before starting any new exercise.</li> <li>• Pace yourself, do not rush. Put off less important activities. Rest when you need to.</li> <li>• Ask family or friends to help you with things like housework, shopping, and child or pet care.</li> <li>• Eat well and drink at least 6 to 8 glasses of water or other liquids every day (unless your health care team has told you to drink more or less).</li> <li>• Avoid driving or using machinery if you are feeling tired.</li> </ul> <p>Ask your health care team for the <a href="#">Fatigue</a> pamphlet for more information.</p>	<p>Talk to your health care team if it does not improve or if it is severe.</p>

<b>Less Common Side Effects (10 to 24 out of 100 people)</b>	
<b>Side effects and what to do</b>	<b>When to contact health care team</b>
<p><b>Constipation</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• Having bowel movements (going poo) less often than normal.</li> <li>• Small hard stools (poo) that look like pellets.</li> <li>• The need to push hard and strain to have any stool (poo) come out.</li> <li>• Stomach ache or cramps.</li> <li>• A bloated belly, feeling of fullness, or discomfort.</li> <li>• Leaking of watery stools (poo).</li> <li>• Lots of gas or burping.</li> <li>• Nausea or vomiting</li> </ul> <p><b>What to do?</b>  <b>To help prevent constipation:</b></p> <ul style="list-style-type: none"> <li>• Try to eat more fiber rich foods like fruits with skin, leafy greens and whole grains.</li> <li>• Drink at least 6 to 8 cups of liquids each day unless your health care team has told you to drink more or less.</li> <li>• Be Active. Exercise can help to keep you regular.</li> <li>• If you take opioid pain medication, ask your health care team if eating more fibre is right for you.</li> </ul> <p><b>To help treat constipation:</b></p> <ul style="list-style-type: none"> <li>• If you have not had a bowel movement in 2 to 3 days you may need to take a laxative (medication to help you poo) to help you have regular bowel movements. Ask your health care team what to do.</li> </ul> <p>Ask your health care team for the <a href="#">Constipation</a> Pamphlet for more information.</p>	<p>Talk to your health care team if it does not improve or if it is severe.</p>
<p><b>Diarrhea</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• Loose, watery, unformed stool (poo) that may happen days to weeks after you get your treatment.</li> </ul> <p><b>What to do?</b>  <b>If you have diarrhea:</b></p> <ul style="list-style-type: none"> <li>• Take anti-diarrhea medication if your health care team prescribed it or told you to take it.</li> </ul>	<p>Talk to your health care team if no improvement after 24 hours of taking diarrhea medication or if severe (more than 7 times in one day).</p>

<b>Less Common Side Effects (10 to 24 out of 100 people)</b>	
<ul style="list-style-type: none"> <li>• Do not eat foods or drinks with artificial sweetener (like chewing gum or 'diet' drinks), coffee and alcohol.</li> <li>• Eat many small meals and snacks instead of 2 or 3 large meals.</li> <li>• Drink at least 6 to 8 cups of liquids each day, unless your health care team has told you to drink more or less.</li> <li>• Talk to your health care team if you can't drink 6 to 8 cups of liquids each day when you have diarrhea. You may need to drink special liquids with salt and sugar, called Oral Rehydration Therapy.</li> <li>• Talk to your health care team if your diarrhea does not improve after 24 hours of taking diarrhea medication or if you have diarrhea more than 7 times in one day.</li> </ul> <p>Ask your health care team for the <a href="#">Diarrhea</a> pamphlet for more information.</p>	
<p><b>Cough and feeling short of breath</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• You may have a cough and feel short of breath.</li> <li>• Symptoms that commonly occur with a cough are: <ul style="list-style-type: none"> <li>◦ Wheezing or a whistling breathing</li> <li>◦ Runny nose</li> <li>◦ Sore throat</li> <li>◦ Heartburn</li> <li>◦ Weight loss</li> <li>◦ Fever and chills</li> </ul> </li> <li>• Rarely this may be severe with chest pain, trouble breathing or coughing up blood.</li> </ul> <p><b>What to do?</b></p> <ul style="list-style-type: none"> <li>• Check your temperature to see if you have a fever. Read the above section <b>“What should I do if I feel unwell, have pain, a headache or a fever?”</b></li> <li>• If you have a fever, try to talk to your health care team. <b>If you are not able to talk to them for advice, you MUST get emergency medical help right away.</b></li> <li>• <b>If you have a severe cough with chest pain, trouble breathing or you are coughing up blood, get medical help right away.</b></li> </ul>	<p>Talk to your health care team. If you are not able to talk to your health care team for advice, and you have a fever or severe symptoms, you <b>MUST</b> get emergency medical help right away.</p>

<b>Less Common Side Effects (10 to 24 out of 100 people)</b>	
<p><b>Mild joint, muscle pain or cramps</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• New pain in your muscles or joints, muscle cramps, or feeling achy.</li> </ul> <p><b>What to do?</b></p> <ul style="list-style-type: none"> <li>• Take pain medication (acetaminophen or opioids such as codeine, morphine, hydromorphone, oxycodone) as prescribed.</li> <li>• Read the above section: <b>“What should I do if I feel unwell, have pain, a headache or a fever?”</b> before taking acetaminophen (Tylenol®), ibuprofen (Advil®, Motrin®), naproxen (Aleve®) or Aspirin. These medications may hide an infection that needs treatment or they may increase your risk of bleeding.</li> <li>• Rest often and try light exercise (such as walking) as it may help.</li> </ul> <p>Ask your health care team for the <a href="#">Pain</a> pamphlet for more information.</p>	<p>Talk to your health care team if it does not improve or if it is severe.</p>
<p><b>Mild swelling</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• You may have mild swelling or puffiness in your arms and/or legs. Rarely, this may be severe.</li> </ul> <p><b>What to do?</b></p> <p><b>To help prevent swelling:</b></p> <ul style="list-style-type: none"> <li>• Eat a low-salt diet.</li> </ul> <p><b>If you have swelling:</b></p> <ul style="list-style-type: none"> <li>• Wear loose-fitting clothing.</li> </ul> <p>For swollen legs or feet, keep your feet up when sitting</p>	<p>Talk to your health care team if it does not improve or if it is severe.</p>
<p><b>Proteins in Urine (pee)</b></p> <p>Your health care team may do urine tests to check for proteins in your pee.</p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• Swelling in your face, legs, or belly.</li> <li>• Recent weight gain that is not normal for you.</li> <li>• Foamy, frothy or bubbly-looking pee.</li> </ul> <p><b>What to do?</b></p> <p>Talk to your healthcare team if it does not improve or if it is severe.</p>	<p>Talk to your health care team if it does not improve or if it is severe.</p>

<b>Less Common Side Effects (10 to 24 out of 100 people)</b>	
<p><b>Nausea and vomiting (generally mild)</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• Nausea is feeling like you need to throw up. You may also feel light-headed.</li> <li>• You may feel nausea within hours to days after your treatment.</li> </ul> <p><b>What to do?</b></p> <p><b>To help prevent nausea:</b></p> <ul style="list-style-type: none"> <li>• It is easier to prevent nausea than to treat it once it happens.</li> <li>• Drink clear liquids and have small meals. Get fresh air and rest.</li> <li>• Do not eat spicy, fried foods or foods with a strong smell.</li> <li>• Limit caffeine (like coffee, tea) and avoid alcohol.</li> </ul> <p><b>If you have nausea or vomiting:</b></p> <ul style="list-style-type: none"> <li>• Take your rescue (as-needed) anti-nausea medication(s) as prescribed.</li> <li>• Ask your health care team for the <a href="#">Nausea and Vomiting</a> pamphlet for more information.</li> <li>• Talk to your health care team if: <ul style="list-style-type: none"> <li>◦ nausea lasts more than 48 hours</li> <li>◦ vomiting lasts more than 24 hours or if it is severe</li> </ul> </li> </ul>	<p>Talk to your health care team if nausea lasts more than 48 hours or vomiting lasts more than 24 hours or if it is severe.</p>
<p><b>Neuropathy (Tingling, numb toes or fingers)</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• Numbness or tingling of your fingers and toes may happen after starting your treatment.</li> <li>• Sometimes it can be painful and feel like burning sensation, which may be severe.</li> </ul> <p><b>What to do?</b></p> <ul style="list-style-type: none"> <li>• <b>Talk to your health care team if you have symptoms of neuropathy.</b></li> <li>• Numbness may slowly get better after your treatment ends.</li> </ul> <p>In rare cases, it may continue long after treatment ends. If you continue to have bothersome symptoms, talk to your health care team for advice.</p>	<p>Talk to your health care team, especially if you have trouble doing tasks like doing up buttons writing, moving, or if you have severe pain or numbness.</p>

<b>Less Common Side Effects (10 to 24 out of 100 people)</b>	
<p><b>Too much or too little salt in your body</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• Muscle spasms, cramping, weakness, twitching, or convulsions.</li> <li>• Irregular heartbeat, confusion or blood pressure changes.</li> </ul> <p><b>What to do?</b></p> <p><b>Get emergency medical help right away for severe symptoms.</b></p>	<p>Get emergency medical help right away for severe symptoms.</p>
<p><b>Low Appetite</b></p> <p><b>What to look for?</b></p> <ul style="list-style-type: none"> <li>• Loss of interest in food or not feeling hungry.</li> <li>• Weight loss.</li> </ul> <p><b>What to do?</b></p> <ul style="list-style-type: none"> <li>• Try to eat your favourite foods</li> <li>• Eat small meals throughout the day.</li> <li>• You may need to take meal supplements to help keep your weight up.</li> <li>• Talk to your health care team if you have no appetite</li> </ul> <p>Ask your health care team for the <a href="#">Loss of Appetite</a> pamphlet for more information.</p>	<p>Talk to your health care team if it does not improve or if it is severe.</p>

**Other rare, but serious side effects are possible.**

**If you experience ANY of the following, speak to your cancer health care provider or get emergency medical help right away:**

- Irregular heartbeat, shortness of breath, chest pain, fainting spells or swelling in your legs, ankles and belly
- Pain, swelling and hardening of the vein in an arm or leg
- Severe chest pain, difficulty breathing or coughing up blood
- Severe rash with blistering or peeling of your skin or any unusual changes in your skin; may also have fever and swollen glands
- Lower back pain, swelling, pee less than usual and have unusual weight gain
- Yellowish skin or eyes, unusually dark pee or pain on the right side of your belly
- Itchiness, rash, swollen lips, face or tongue, chest and throat tightness
- Unusual weight gain with some of the following: feeling tired or having low energy, dry skin, nails or hair that breaks easily, and feeling sensitive to coldness

